

ISSN 1859 - 4735



JOURNAL OF MEDICINAL MATERIALS

No. 1+2 - Vol. 31
3/2026

SPECIAL ISSUE

THE 14th SCIENTIFIC CONFERENCE OF NATIONAL INSTITUTE OF MEDICINAL MATERIALS (NIMM)
THE 65th ANNIVERSARY OF NIMM FOUNDATION (13/4/1961-13/4/2026)

NATIONAL INSTITUTE OF MEDICINAL MATERIALS

3B QUANG TRUNG - CUA NAM - HANOI - VIETNAM
TEL: (+84-24) 38247058 FAX: (+84-24) 39348740
Email: tapchiduoclieu@nimm.org.vn; tapchiduoclieu@gmail.com

EDITOR-IN-CHIEF

Assoc. Prof. PhD. Do Thi Ha

I. Subcommittee of Pharmacognosy - Traditional Medicine

• **Assoc. Prof. Dr. Sci. Nguyen Minh Khoi - Head of Subcommittee**

- Assoc. Prof. PhD. Nguyen Thuong Dong - Member
- Prof. PhD. Nguyen Minh Duc - Member
- Prof. PhD. Tran Cong Luan - Member
- Prof. PhD. Won Keun Oh - Member
- Assoc. Prof. PhD. Tran Hung - Member
- Assoc. Prof. PhD. Nguyen Thi Bich Thu - Member
- Assoc. Prof. PhD. Phuong Thien Thuong - Member
- Assoc. Prof. PhD. Phan Minh Giang - Member
- Assoc. Prof. PhD. Le Nguyen Thanh - Member
- Assoc. Prof. PhD. Le Viet Dung - Member
- Assoc. Prof. PhD. Nguyen Quoc Huy - Member
- Assoc. Prof. PhD. Nguyen Thu Hang - Member
- PhD. Tran Minh Ngoc - Member
- PhD. Nguyen Van Tai - Member
- Assoc. Prof. PhD. Le Dinh Chi - Member
- PhD. Nguyen Tuan Hiep - Member.

II. Subcommittee of Pharmacology - Clinical Pharmacy

• **Assoc. Prof. PhD. Pham Thi Nguyet Hang - Head of Subcommittee**

- Assoc. Prof. Dr.Sci. Do Trung Dam - Member
- Assoc. Prof. PhD. Nguyen Thi Thu Huong - Member
- Assoc. PhD. William R. Folk - Member
- Assoc. Prof. PhD. Pham Thi Van Anh - Member
- Assoc. Prof. PhD. Nguyen Thuy Duong - Member
- Assoc. Prof. PhD. Le Van Minh - Member
- PhD. Tran Thi Hien - Member.

III. Subcommittee of Agriculture - Biology

• **PhD. Nguyen Van Khiem - Head of Subcommittee**

- PhD. Phan Thuy Hien - Member
- Assoc. Prof. PhD. Nguyen Van Tap - Member
- Assoc. Prof. PhD. Pham Thanh Huyen - Member
- Prof. PhD. Tran The Bach - Member.

IV. Subcommittee of Secretariat Administration

• **Assoc. Prof. PhD. Le Nguyen Thanh - Head of Subcommittee**

- PhD. Le Thi Xoan - Academic Secretary
- BSc. Nguyen Thi Minh Loc - Administrative Secretary.

CONTENTS VOL.31(1+2), 2026 (SPECIAL ISSUE)
(Pages 1 - 160)

Page	SCIENTIFIC RESEARCH
3	Foreword: Special Issue Commemorating the 65th Anniversary of the National Institute of Medicinal Materials and Welcoming the 1st International Conference V.IMPACT-2026 - Assoc. Prof. Do Thi Ha, PhD
	REVIEW
4	Results of Scientific Activities of the National Institute of Medicinal Materials (2016–2025) and Development Orientation Toward 2030 - Tran Minh Ngoc, Do Thi Ha, Pham Tuan Anh
	SCIENTIFIC RESEARCH
10	Triterpenoid Constituents from the Leaves of <i>Walsura pinnata</i> Hassk - Bui Huu Tai, Nguyen Viet Dung, Nguyen Huy Hoang, Ngo Anh Bang, Duong Thi Dung, Duong Thi Hai Yen, Phan Thi Thanh Huong, Dan Thuy Hang, Pham Hai Yen, Phan Van Kiem, Nguyen Phuong Thao
17	Chemical Constituents from the Propolis of the Stingless Bee <i>Lepidotrigona terminata</i> (Smith, 1878) Collected in Gia Lai - Diep Thi Lan Phuong, Nguyen Thu Thao Van, Vo Thi Thanh Tuyen, Nguyen Thi Nghia, Pham Ngoc Thach, Le Thi Cam Nhung, Le Duy Thanh, Vo Manh Tien, Tran Minh Ngoc, Le Nguyen Thanh
23	Glycoside Constituents from <i>Abroma augustum</i> and their <i>in vitro</i> Antiproliferative Activity - Ninh Thi Ngoc, Pham Thi Cham, Le Thi Vien, Tran Thi Hong Hanh, Do Thi Thao, Nguyen Xuan Cuong, Nguyen Hoai Nam
29	Chemical Profiling and Anti-Inflammatory Effects of <i>Physalis angulata</i> in Lipopolysaccharide-Stimulated Bone Marrow–Derived Dendritic Cells - Dinh Thi Huong, Le Ba Vinh, Nguyen Van Chinh, Nguyen Cao Cuong
36	Phenolic Constituents from the Ethyl Ecetate Extract of <i>Camellia flava</i> Flowers (Theaceae) Collected in Phu Tho Province - Do Thi Thuy, Nguyen Duong Hong Anh, Tran The Bach, Do Van Hieu, Do Duc Nhuan, Than Kieu My, Tran Minh Ngoc, Nguyen Thanh Cong, Le Nguyen Thanh
42	Lignans and Other Compounds from the Leaves of <i>Triadica rotundifolia</i> with Inhibitory Activity on Nitric Oxide Production - Ninh Khac Ban, Tran My Linh, Ninh Khac Thanh Tung, Bui Van Thanh, Pham Hai Yen, Bui Huu Tai, Phan Van Kiem, Nguyen Phuong Thao
49	Chemical Constituents of <i>Ganoderma multipileum</i> Ding Hou, Ganodermataceae - Nguyen Thi Duyen, Tran Thi Tuyet, Phan Quynh Trang, Nguyen Trung Hieu, Nguyen Duc Hieu, Nguyen Phuong Dai Nguyen
53	Chemical Composition, Antimicrobial and Cytotoxic Activities of the Leaf Essential Oil of <i>Knema globularia</i> (Lamk.) Warb. from Thanh Hoa, Vietnam - Luong Le Uyen Trang, Pham Thi Huong Ly, Nguyen Thi Hoa Hien, Nguyen Thanh Huyen, Nguyen Thi Hien, Nguyen Thi Thu, Khong Thi Hoa

60	Azaphilone Derivatives Isolated from Endophytic Fungus <i>Penicillium sclerotiorum</i> - Ly Tu Loan, Vo Van Leo, Ma Chi Thanh
67	Chemical Constituents from the Aerial Parts of <i>Barleria lupulina</i> (Acanthaceae) - Ngo Thi Ngoc Yen, Do Hoang Anh, Nguyen Thi Thu, Nguyen Thi Ha Ly, Le Nguyen Thanh, Nguyen Quynh Nga, Dao Viet Quoc, Nguyen Thu Hang, Nguyen Hoang Minh, Nguyen Minh Khoi, Le Tran Nguyen Vu
74	Determination of Essential Oil Content, Composition, and Total Polyphenol Content of <i>Rhizoma homalomenae occulta</i> Collected in Vietnam - Nguyen Thi Ha Ly, Pham Thi Hien, Ngo Thi Thanh Huong, Hoang Dieu Linh, Tran Huyen Trang, Nguyen Minh Khoi
80	Establishment of an In-House Standard for <i>Prunellae spica</i> - Tran Huyen Trang, Pham Thi Hien, Nguyen Thi Ha Ly, Phan Van Truong, Vu Thi Diep, Nguyen Tra My, Nguyen Thi Hang, Nguyen Duong Hong Anh, Do Thi Ha
88	HPTLC Scanner Method for the Simultaneous Determination of Lycopene and β-Carotene in Gac Seed Aril - Pham Thi Hien, Pham Thi Khuyen, Nguyen Thi Hang, Nguyen Thi Lan Hoa, Ta Thi Thao, Do Thi Ha, Nguyen Thi Ha Ly
96	Optimization of the Extraction Process from Yellow Camellia Flowers (<i>Camellia cucphuongensis</i> Ninh & Rosmann) Collected in Phu Tho Province Using Response Surface Methodology - Nguyen Hoang Viet, Nguyen Dinh Quan, Do Duc Nhuan, Nguyen Duong Hong Anh, Tran Thi Mai Anh, Doan Van Nghia, Vu Xuan Hoa, Dang Mai Thien Ha, Nguyen Thanh Cong
105	Antioxidant, Antihyperglycemic, and Renal Protective Effects of <i>Hibiscus tiliaceus</i> L. Leaf Extract on Streptozotocin-Induced Hyperglycemia - Nguyen Thi Thu Huong, Nhieu Chinh Hoang, Nguyen Hoang Minh
113	Acute Oral Toxicity and Laxative Effects of <i>Pharbitis nil</i> (L.) Choisy Seed Extract - Nguyen Thi Kim Oanh, Tran Vien Thong, Truong Huy Hieu, Ly Hai Trieu, Pham Thi Nguyet Hang, Nguyen Thi Vinh Hue, Do Thi Ha
123	Evaluation of Memory-Enhancing Effects of <i>Morinda officinalis</i> How. Root Extracts in a Transgenic <i>hAPP Drosophila</i> Model of Alzheimer's Disease - Nguyen Van Hiep, Pham Thi Nguyet Hang, Trinh Dinh Luc, Leu Khanh Duy, Pham Anh Tung, Nguyen Thi Phuong, Nguyen Van Tai
131	Anti-Inflammatory Effects of the Bark of <i>Phellodendron amurense</i> Rupr. in Experimental Models - Do Thi Hang, Le Thi Kim Van, Do Van Dung, Bui Thanh Tung
138	Neuroprotective Effects of Tianma Gouteng Decoction in a Middle Cerebral Artery Occlusion Model of Ischemic Stroke - Le Thi Xoan, Trinh Lan Phuong, Nguyen Van Hiep, Pham Thi Nguyet Hang, Nguyen Van Tai, Nguyen Thi Thu Trang, Nguyen Thi Thanh Loan, Nguyen Hung Son
147	Investigation of the Variation of Gymnemagenin Content in <i>Gymnema R.Br</i> Under the Influence of Certain Strains of Bacteria - Tong Thi Thanh Vuong, Tran Thi Minh Thuy, Le Thi Kim Van
154	Comparative Transcriptome Profiling of <i>Momordica cochinchinensis</i> (Lour.) Spreng. Aril Using Oxford Nanopore and Illumina Sequencing Technologies - Tran Duc Trung, Nguyen Hoang, Le Thi Tu Linh, Nguyen Thi Thu Hang, Pham Thanh Huyen, Truong Nhat My, Nguyen Thi Lan Hoa

TRITERPENOID CONSTITUENTS FROM THE LEAVES OF *WALSURA PINNATA* HASSK

Bui Huu Tai, Nguyen Viet Dung, Nguyen Huy Hoang, Ngo Anh Bang,
Duong Thi Dung, Duong Thi Hai Yen, Phan Thi Thanh Huong, Dan Thuy Hang,
Pham Hai Yen, Phan Van Kiem, Nguyen Phuong Thao*
Institute of Chemistry, Vietnam Academy of Science and Technology (VAST),
Hanoi, Vietnam

*Corresponding author: thaonp@ich.vast.vn

Received December 9th, 2025

Accepted January 8th, 2026

Summary

By employing chromatographic techniques, six triterpenoids (**1–6**) were successfully isolated from the leaves of *Walsura pinnata* Hassk. Their structures were elucidated to be cumingianol E (**1**), 21-*O*-methyltoosendanpentol (**2**), 21 α -methylmelianodiol (**3**), 21*S*,23*R*-epoxy-3 β ,24*R*-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-en (**4**), odoratone (**5**), and cabraleone (**6**) by detailed analysis of 1D, 2D NMR, and MS spectroscopic data, as well as comparison with previously reported literature. This is the first time compounds **1–6** from *W. pinnata* were reported.

Keywords: *Walsura pinnata*; Meliaceae; Apotirucallane; Tirucallane; Dammarane; Triterpenoids.

1. Introduction

Plants of the genus *Walsura* Roxb, belonging to the Meliaceae family and comprising approximately 21 species [1], are distributed in subtropical regions, including South India, mainland China, and Southeast Asia (Malaysia, Indonesia, Thailand, and Vietnam) [2]. Currently, there are no accurate data regarding the exact number of *Walsura* species present in Vietnam. However, based on published literature, seven species have been recorded in the country: *W. bonii*, *W. cochinchinensis*, *W. elata*, *W. pinnata*, *W. poilanei*, *W. robusta*, and *W. villosa* [3]. Traditionally, members of this genus have been used in folk medicine to treat various ailments, such as tendon disabilities, hemorrhoids, excessive bleeding, and wounds, as well as for their expectorant, emmenagogue, and emetic properties [4]. Previous phytochemical investigations on *Walsura* have led to the isolation of phenols, limonoids, lignans, and terpenoids, some of which exhibited cytotoxic, antimicrobial, antioxidant, antifeedant, and neuroprotective activities [5],[6],[7],[8]. However, phytochemical and pharmacological data on *W. pinnata* remain limited [4]. To date, a few studies have reported the presence of sesquiterpenoid and triterpenoid constituents, which have drawn interest due to their cytotoxic effects [9],[10],[11]. In continuation of our interest in the phytochemistry of plants belonging to the genus *Walsura*, we have conducted further

investigations and successfully isolated six triterpenoid compounds (**1–6**, Fig. 1) from the leaves of *W. pinnata*.




Fig. 1. Chemical structures of compounds **1–6**

2. Materials and methods

2.1. Chemicals and equipment

Optical rotations were determined on a JASCO P-2000 polarimeter. The NMR spectra were recorded on a Bruker AVANCE NEO 600 FT-NMR spectrometer with TMS as an internal standard. The HRESIMS data were taken on an Agilent 6530 Accurate Mass Q-TOF LC/MS. Column chromatography (CC) was conducted on silica gel, Diaion HP-20, Sephadex LH-20, and RP-18. Preparative HPLC was performed using Agilent 1100, J' sphere ODS H-80 column (4

μm , $250 \times 20 \text{ mm}$), a flow rate of 6 mL/min, and a detector DAD at wavelengths of 205, 230, 254, and 280 nm. An isocratic mobile phase with a flow rate of 3.0 mL/min was used in pre-HPLC. Analytical thin-layer chromatography was performed on precoated *silica gel* 60 F₂₅₄ and RP-18 F₂₅₄S plates, and compounds were visualized by spraying with 10% H₂SO₄ in water and then heating for 1.5–2 minutes.

2.2. Plant material

The leaves of *W. pinnata* were collected at Dakrong, Quang Tri Province, Vietnam, in May 2025. The species was taxonomically identified by Dr. Le Tuan Anh (Vietnam National Museum of Nature). A voucher herbarium specimen (NCCT-P184) has been deposited at the Herbarium of the Institute of Chemistry, VAST.

2.3. Extraction and isolation

The dried leaves of *W. pinnata* (3.5 kg) were ultrasonically extracted with methanol ($3 \times 7.0 \text{ L}$, each for 2.5 h at room temperature), yielding a MeOH extract (300 g) after solvent removal under reduced pressure. The extract was then suspended in water and sequentially partitioned with organic solvents to obtain the *n*-hexane (H, 32.5 g), dichloromethane (D, 95 g), ethyl acetate (E, 40 g), and aqueous fractions. HPLC analysis indicated that the CH₂Cl₂ fraction contained a relatively high abundance of secondary metabolites, and this fraction was therefore selected for further chromatographic separation. The CH₂Cl₂ fraction (95 g) was subjected to *silica gel* CC, using a gradient solvent system of CH₂Cl₂-MeOH (from 50:0 to 0:100), based on TLC analysis results. Fractions with similar polarity were combined, resulting in the acquisition of seven fractions (D1–D7). Fraction D2 (5.1 g) was purified on an RP-18 column using 40% MeOH in H₂O as the eluent, yielding six subfractions (D2.1–D2.6). Among these, subfraction D2.2 (56 mg) was further purified by semi-preparative HPLC with CH₃CN-H₂O (65:35) as the mobile phase, yielding compound **1** (6.5 mg, $t_R = 44.3 \text{ min}$). Similarly, subfraction D2.3 (82 mg) was purified by a semi-preparative HPLC system using a CH₃CN-H₂O (50:50) mixture, resulting in the isolation of compound **2** (10.8 mg, $t_R = 33.8 \text{ min}$). Subfraction D3 (1.52 g) was subjected to RP-18 CC with MeOH-H₂O (2:3) as the eluent, producing three subfractions (D3.1–D3.3). Further purification of subfraction D3.2 (75 mg) by semi-preparative HPLC, using CH₃CN-H₂O (30:70) as the mobile phase, yielded compound **6** (7.2 mg, $t_R = 47.3 \text{ min}$).

Additionally, subfraction D3.3 (0.8 g) was subjected to *silica gel* CC using a CH₂Cl₂-MeOH-H₂O solvent system (5:1:0.1, v/v/v), yielding five fractions (D3.3a to D3.3e). Compounds **3** (10.1 mg, $t_R = 31.8 \text{ min}$) and **5** (7.9 mg, $t_R = 35.4 \text{ min}$) were obtained from subfraction D3.3b (0.11 g) through semi-preparative HPLC system (CH₃CN-H₂O containing 0.05% TFA, 20:80). In a similar process to that described above, subfraction D3.3d (0.10 g) was purified by semi-preparative HPLC using 20% CH₃CN in H₂O as the mobile phase, yielding compound **4** (8.9 mg, $t_R = 40.5 \text{ min}$).

Cumingianol E (1): White amorphous powder; $[\alpha]_D^{23}$: -89.5 (c 0.1, CHCl₃); HRESIMS m/z 557.3840 $[M + Na]^+$ (calcd. for $[C_{32}H_{54}NaO_6]^+$, 557.3835); ¹H (600 MHz, CDCl₃) and ¹³C NMR (150 MHz, CDCl₃) spectroscopic data, see Table 1.

21-O-Methyltoosendanpentol (2): White amorphous powder; $[\alpha]_D^{23}$: -41.4 (c 0.1, CHCl₃); HRESIMS m/z 543.3679 $[M + Na]^+$ (calcd. for $[C_{31}H_{52}NaO_6]^+$, 543.3676); ¹H (600 MHz, CDCl₃) and ¹³C NMR (150 MHz, CDCl₃) spectroscopic data, see Table 1.

21a-Methylmelianodiol (3): White amorphous powder; $[\alpha]_D^{23}$: -89.7 (c 0.1, CHCl₃); HRESIMS m/z 503.3739 $[M + H]^+$ (calcd. for $[C_{31}H_{51}O_5]^+$, 503.3731); ¹H (600 MHz, CDCl₃) and ¹³C NMR (150 MHz, CDCl₃) spectroscopic data, see Table 2.

21S,23R-epoxy-3 β ,24R-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-en (4): White amorphous powder; $[\alpha]_D^{23}$: $+35.0$ (c 0.1, CHCl₃); HRESIMS m/z 541.3874 $[M + Na]^+$ (calcd. for $[C_{32}H_{54}NaO_5]^+$, 541.3864); ¹H (600 MHz, CDCl₃) and ¹³C NMR (150 MHz, CDCl₃) spectroscopic data, see Table 2.

Odoratone (5): Colorless needles; $[\alpha]_D^{23}$: -41.8 (c 0.1, CHCl₃); ¹H NMR (CDCl₃, 600 MHz) δ (ppm): 1.45 (1H, m, H_a-1), 1.98 (1H, m, H_b-1), 2.24 (1H, dd, $J = 14.4, 1.6 \text{ Hz}$, H_a-2), 2.75 (1H, ddd, $J = 14.4, 14.4, 5.4 \text{ Hz}$, H_b-2), 1.72 (1H, t, $J = 9.0 \text{ Hz}$, H-5), 2.10 (2H, m, H-6), 5.31 (1H, dd, $J = 6.6, 3.6 \text{ Hz}$, H-7), 2.28 (1H, m, H-9), 1.56 (2H, m, H-11), 1.62 (1H, m, H_a-12), 1.86 (1H, m, H_b-12), 1.50 (2H, m, H-15), 1.37 (1H, m, H_a-16), 2.02 (1H, m, H_b-16), 1.87 (1H, m, H-17), 0.83 (3H, s, H-18), 1.00 (3H, s, H-19), 1.64 (1H, m, H-20), 0.85 (3H, d, $J = 7.2 \text{ Hz}$, H-21), 3.82 (1H, dd, $J = 6.6, 1.8 \text{ Hz}$, H-22), 3.97 (1H, dd, $J = 6.0, 5.4 \text{ Hz}$, H-23), 3.66 (1H, dd, $J = 6.0, 5.4 \text{ Hz}$, H-24), 1.22 (3H, s, H-26), 1.22 (3H, s, H-27), 1.05 (3H, s, H-28), 1.11 (3H, s, H-29), and 1.03 (3H, s, H-30); ¹³C NMR (CDCl₃, 150 MHz) δ (ppm):

38.6 (C-1), 35.1 (C-2), 217.1 (C-3), 47.9 (C-4), 52.4 (C-5), 24.4 (C-6), 117.8 (C-7), 146.0 (C-8), 48.6 (C-9), 35.0 (C-10), 18.3 (C-11), 33.6 (C-12), 43.6 (C-13), 51.2 (C-14), 34.1 (C-15), 27.9 (C-16), 49.3 (C-17), 21.9 (C-18), 12.8 (C-19), 37.5 (C-20), 12.4 (C-21), 83.7 (C-22), 72.9 (C-23), 77.4 (C-24), 80.9 (C-25), 27.7 (C-26), 21.3 (C-27), 24.6 (C-28), 21.6 (C-29), and 27.7 (C-30).

Cabraleone (6): White amorphous powder; $[\alpha]_D^{23}$: +67.3 (*c* 0.1, CHCl₃); ¹H NMR (CDCl₃, 600 MHz) δ (ppm): 1.45 (1H, m, H_a-1), 1.92 (1H, m, H_b-1), 2.43 (1H, m, H_a-2), 2.50 (1H, m, H_b-2), 1.37 (1H, dt, *J* = 9.0, 3.0 Hz, H-5), 1.47 (1H, m, H_a-6), 1.55 (1H, m, H_b-6), 1.32 (1H, m, H_a-7), 1.56 (1H, m, H_b-7), 1.43 (1H, m, H-9), 1.25 (1H, m, H_a-11), 1.52 (1H, m, H_b-11), 1.80 (1H, m, H_a-12), 1.88 (1H, m, H_b-12), 1.60 (1H, m, H-13), 1.10 (1H, m, H_a-15), 1.47 (1H, m, H_b-15), 1.25 (1H, m, H_a-16), 1.89 (1H, m, H_b-16), 1.82 (1H, m, H-17), 0.99 (3H, s, H-18), 0.94 (3H, s, H-19), 1.14 (3H, s, H-21), 1.63 (1H, m, H_a-22), 1.72 (1H, m, H_b-22), 1.48 (1H, m, H_a-23), 1.78 (1H, m, H_b-23), 3.73 (1H, dd, *J* = 7.8, 7.2 Hz, H-24), 1.21 (3H, s, H-26), 1.12 (3H, s, H-27), 1.08 (3H, s, H-28), 1.04 (3H, s, H-29), and 0.88 (3H, s, H-30); ¹³C NMR (CDCl₃, 150 MHz) δ (ppm): 39.9 (C-1), 34.1 (C-2), 218.1 (C-3), 47.4 (C-4), 55.3 (C-5), 19.7 (C-6), 34.6 (C-7), 40.3 (C-8), 50.2 (C-9), 36.9 (C-10), 22.1 (C-11), 26.1 (C-12), 43.1 (C-13), 50.1 (C-14), 31.4 (C-15), 27.5 (C-16), 49.5 (C-17), 15.1 (C-18), 16.0 (C-19), 86.3 (C-20), 23.6 (C-21), 35.7 (C-22), 25.7 (C-23), 83.3 (C-24), 71.4 (C-25), 27.4 (C-26), 24.3 (C-27), 26.7 (C-28), 21.0 (C-29), and 16.4 (C-30).

3. Results and discussion

Compound **1** was obtained as a white amorphous powder. Its molecular formula was determined to be C₃₂H₅₄O₆, corresponding to six degrees of unsaturation, based on its sodium adduct molecular ion peak at *m/z* 557.3840 [*M* + Na]⁺ (calcd. for [C₃₂H₅₄NaO₆]⁺, 557.3835) in the HRESIMS spectrum. The ¹H NMR spectrum of **1** exhibited signals for seven tertiary methyl groups [δ_H 0.85, 0.90, 1.05, 1.06, 1.15, and 1.24 (3H, s, for each)], five oxymethine protons [δ_H 3.40 (1H, br s, H-3), 3.90 (1H, t, *J* = 3.0 Hz, H-7), 4.80 (1H, d, *J* = 3.0 Hz, H-21), 4.21 (1H, ddd, *J* = 10.8, 4.8, 2.4 Hz, H-23), and 3.37 (1H, d, *J* = 2.4 Hz, H-24)], two methoxy groups (δ_H 3.23 and 3.35), and a characteristic olefinic proton signal [δ_H 5.42 (1H, d, *J* = 1.8 Hz, H-15)]. All 32 carbons in the molecular structure were resolved as individual resonances in the ¹³C NMR spectrum and classified using HSQC spectroscopy as two

olefinic carbons [δ_C 162.4 (C-14) and 119.1 (C-15)], one acetal carbon [δ_C 109.2 (C-21)], four *sp*³ quaternary carbons [δ_C 37.0 (C-4), 44.4 (C-8), 37.7 (C-10), and 47.0 (C-13)], and one oxygen-bearing *sp*³ quaternary carbon [δ_C 77.0 (C-25)] (Table 1). All these data, coupled with a literature survey, indicated that **1** is an apotirucallane-type triterpenoid and shares the same A–D ring framework as cumingianol E [12]. Four rings and one double bond account for five of the six degrees of unsaturation, indicating that the remaining degree of unsaturation must arise from an additional ring. A spin system, CH(17)–CH(20)–CH(21)–CH(24)–CH(23)–CH₂(22)–CH(20), determined from the ¹H–¹H COSY spectrum (Fig. 2), suggested that C-21 is connected to C-20, C-22, and C-23. This connectivity requires the presence of an additional five-membered ring (ring E) attached at C-17. This structural feature was confirmed by HMBC correlations of H-21 (δ_H 4.80) with C-17, C-20, C-22, and C-23 (δ_C 57.7, 46.2, 35.5, and 75.1, respectively) as well as correlations of H_a-22 (δ_H 1.60) and H_b-22 (δ_H 1.97) with C-17, C-20, C-23, and C-24 (δ_C 76.3). HMBC correlations from the proton signals at δ_H 1.15 (H₃-26) and 1.24 (H₃-27) to the carbon signals at δ_C 76.3 (C-24) and 77.0 (C-25), together with the methoxy signal (δ_H 3.23) correlating with C-25, supported the presence of an isopropyl methoxy group at C-25. In addition, HMBC cross-peaks between H-21 and C-23, and between a methoxyl signal (δ_H 3.35) and C-21, indicated a tetrahydrofuran moiety bearing a methoxy group at C-21. Furthermore, HMBC correlations of H₃-28 (δ_H 0.95) with C-3, C-4, C-5, and C-29, and of H₃-30 (δ_H 1.05) with C-7, C-8, C-9, and C-14 were consistent with the presence of OH-3, OH-7, and $\Delta^{14,15}$ substituents, respectively. The hydroxy groups at C-3 and C-7 were assigned as α -oriented based on the broad singlet of H-3 and the small coupling constant values of H-7 (*J* = 3.0 Hz) [12].




Fig. 2. Key COSY and HMBC correlations of **1** and **3**

Table 1. NMR spectroscopic data of compounds **1** and **2** and the reference compounds

No.	# δ_C	1		δ_C^a	2	
		$\delta_C^{a,b}$	$\delta_H^{a,c}$ mult. (J in Hz)		$\delta_C^{a,b}$	$\delta_H^{a,c}$ mult. (J in Hz)
1	33.3	32.6	1.33 m/1.38 m	32.8	32.5	1.35 m/1.38 m
2	26.1	25.0	1.55 m/1.92 m	25.0	25.0	1.57 m/1.92 m
3	74.9	76.2	3.40 br s	76.2	76.1	3.39 br s
4	37.5	37.0	-	37.1	37.0	-
5	40.6	40.4	1.98 m	40.5	40.3	1.98 m
6	25.0	23.7	1.71 m	23.7	23.6	1.71 m
7	72.5	72.3	3.90 t (3.0)	72.3	72.3	3.91 br s
8	44.4	44.4	-	44.3	44.3	-
9	42.0	41.6	2.02 dd (12.0, 7.8)	41.6	41.6	2.02 dd (11.4, 7.2)
10	38.0	37.7	-	37.7	37.6	-
11	16.6	16.2	1.50 m/1.74 m	16.3	16.2	1.50 m/1.72 m
12	33.3	32.7	1.43 m/1.60 m	32.6	32.7	1.45 m/1.79 m
13	47.0	47.0	-	46.9	46.9	-
14	162.2	162.4	-	162.4	162.3	-
15	118.7	119.1	5.42 d (1.8)	119.2	119.1	5.43 br s
16	34.9	34.7	2.10 m/2.16 m	34.7	34.6	2.11 m/2.19 m
17	58.0	57.7	1.71 m	57.5	57.4	1.92 m
18	19.2	19.3	1.06 s	19.3	19.2	1.06 s
19	15.6	15.2	0.90 s	15.2	15.2	0.90 s
20	46.3	46.2	2.33 m	45.9	45.5	2.33 m
21	109.5	109.2	4.80 d (3.6)	109.6	109.4	4.80 d (3.0)
22	35.2	35.5	1.60 dd (12.0, 10.8) 1.97 ddd (12.0, 7.2, 4.8)	33.8	33.8	1.78 dd (12.0, 10.2) 1.95 ddd (12.0, 8.4, 4.8)
23	68.9	75.1	4.21 ddd (10.8, 4.8, 2.4)	76.9	76.7	4.24 ddd (10.2, 8.4, 4.8)
24	76.4	76.3	3.37 d (2.4)	75.6	75.6	3.24 d (4.8)
25	76.4	77.0	-	73.1	73.0	-
26	22.8	20.1	1.15 s	26.4	26.2	1.26 s
27	20.3	21.6	1.24 s	26.5	26.4	1.29 s
28	29.0	28.0	0.95 s	28.1	28.0	0.95 s
29	22.5	22.1	0.85 s	22.1	22.1	0.85 s
30	28.3	27.7	1.05 s	27.7	27.6	1.05 s
21-OCH ₃	55.0	55.4	3.35 s	55.6	55.5	3.35 s
25-OCH ₃	49.0	49.1	3.23 s			

Measured in $^a\text{CDCl}_3$, b150 MHz, and c600 MHz. $^{\#}\delta_C$ of cumingianol *E* in pyridine- d_5 + D_2O [12] and $^{\$}\delta_C$ of 21-O-methyltoosendanpentol in CDCl_3 [13].

Moreover, the *S*-configuration at C-24 was confirmed on the basis of the small coupling constant ($J_{\text{H-23/H-24}} = 2.4$ Hz) in the ^1H NMR spectrum [12],[13]. The NOESY correlations of H_3 -29/ H -2 β , H -2 β / H_3 -19, H_3 -19/ H_3 -30, and H_3 -30/ H -17 indicated that these methyl groups are cofacial and β -oriented, whereas the correlations of H_3 -28/ H -5 and H -9/ H_3 -18 supported an α -orientation. Additionally, the cross-peak observed for H_3 -18/ H -20, H_a -22/ H -23, and H -23/ H_3 -26 showed that both H -20 and H -23 are α -oriented, whereas those for H -17/ H -21, H -21/ H -24 revealed that H -21 and H -24 are β -oriented. Based on the above evidence and comparison with the published data [12], compound **1** was finally identified as cumingianol *E* (Fig. 1 and Table 1). This compound was previously obtained from the leaves of *Dysoxylum cumingianum* (Meliaceae) and was reported to show no growth inhibitory effects against KB, MCF-7, and KB-C2 cell lines [12].

Compound **2** was isolated as a white

amorphous powder. The HRESIMS data of **2** showed a sodium adduct ion at m/z 543.3679 [$\text{M} + \text{Na}$] $^+$ (calcd. for $[\text{C}_{31}\text{H}_{52}\text{NaO}_6]^+$, 543.3676), consistent with the molecular formula $\text{C}_{31}\text{H}_{52}\text{O}_6$ (six degrees of unsaturation). Analysis of the NMR spectra allowed the assignment of most functional groups characteristic of an apotirucallane-type triterpenoid skeleton [12],[13]. Complete elucidation of 1D NMR (Table 1) and HSQC spectra indicated that the structure of **2** closely resembled that of cumingianol *E* (**1**) [12] (Table 1), except for minor differences observed at C-25 resonance in the side chain. The difference between the two compounds is that the methoxy group at C-25 in **1** was replaced by a hydroxy group in **2**. This substitution was suggested by the upfield shifts of C-25, from δ_C 77.0 in **1** to δ_C 73.0 in **2** (Table 1). This conclusion was further supported by 2D HMBC correlations from the proton signal at δ_H 1.26 (H_3 -26)/1.29 (H_3 -27) with C-25 (δ_C 73.0), as well as a 14 mass units difference between the

two molecules. The structure of **2** was additionally confirmed through 2D NMR experiments, including HSQC, HMBC, and NOESY spectra. Its relative configuration of **2** was inferred from NOESY correlations and found to be identical to that of compound **1**. Detailed examination of the NMR spectroscopic data supported the identification of **2** and in comparison with reported literature, established it as 21-*O*-methyltoosendanpentol [14]. This compound was previously reported from the leaves and stems of *Aglaia duperreana* [14] and the fruits of *Melia toosendan* [15]. Thus far, no activity was found in an inhibition assay for compound **2**.

Compound **3** was obtained as a white amorphous powder. Its molecular formula, C₃₁H₅₀O₅, was established based on the HRESIMS peak at *m/z* 503.3739 [M + H]⁺ (calcd. for [C₃₁H₅₁O₅]⁺, 503.3731) and NMR data, indicating seven degrees of unsaturation. The ¹H NMR spectrum of **3** showed signals corresponding to seven methyl groups, one

methoxy group [δ_{H} 3.34 (3H, s)], three oxygenated methine protons, one olefinic proton [δ_{H} 5.32 (1H, dd, *J* = 6.6, 3.0 Hz, H-7)], and several overlapping resonances for aliphatic methine and methylene protons. Combined analysis of the ¹³C NMR and HSQC spectra revealed 31 carbon signals attributable to one ketone group, seven methyl groups, one methoxy carbon, eight *sp*³ methylenes, seven *sp*³ methines (three oxygenated at δ_{C} 109.0, 76.8, and 75.6), one *sp*² methine carbon (δ_{C} 118.2), one *sp*² non-protonated carbon (δ_{C} 145.6), and five *sp*³ non-protonated carbons (Table 2). The remaining five degrees of unsaturation indicated that **3** possessed a pentacyclic core. The presence of a tetrahydrofuran ring in the side chain was assigned based on correlations observed in the 2D NMR data. Overall, the NMR spectroscopic data closely matched those reported for a 3-oxotirucalla-7-ene skeleton bearing a 21,23-epoxy-21-methoxy-24,25-dihydroxy side chain [16].

Table 2. NMR spectroscopic data of compounds **3** and **4** and the reference compounds

No.	# $\delta_{\text{C}}^{\text{a}}$	3		$\delta_{\text{C}}^{\text{a}}$	4	
		$\delta_{\text{C}}^{\text{a,b}}$	$\delta_{\text{H}}^{\text{a,c}}$ mult. (<i>J</i> in Hz)		$\delta_{\text{C}}^{\text{a,b}}$	$\delta_{\text{H}}^{\text{a,c}}$ mult. (<i>J</i> in Hz)
1	38.5	38.5	1.47 m/1.98 m	37.2	37.2	1.14 m/1.67 m
2	35.1	35.1	2.24 dt (14.4, 3.6) 2.75 ddd (14.4, 14.4, 5.4)	27.7	27.7	1.60 m 1.64 m
3	216.9	216.8	-	79.2	79.2	3.23 m
4	47.9	47.9	-	38.9	39.0	-
5	52.3	52.4	1.72 dd (9.6, 7.8)	50.7	50.8	1.31 m
6	24.3	24.4	2.09 m/2.11 m	24.0	24.0	1.97 m/2.14 m
7	118.1	118.2	5.32 dd (6.6, 3.0)	118.2	118.2	5.26 q (3.0)
8	145.5	145.6	-	145.5	145.6	-
9	48.3	48.4	2.33 m	48.8	48.9	2.24 m
10	34.9	34.9	-	35.0	35.1	-
11	17.7	17.8	1.59 m	17.7	17.7	1.52 m
12	31.5	31.6	1.55 m/1.75 m	31.2	31.2	1.30 m/1.87 m
13	43.60	43.8	-	43.5	43.6	-
14	50.9	51.0	-	50.7	50.8	-
15	34.3	34.4	5.48 d (1.8)	34.2	34.2	1.49 m/1.54 m
16	27.4	27.4	1.75 m/1.95 m	27.3	27.3	1.29 m/1.87 m
17	50.3	50.3	1.30 m/1.94 m	45.0	45.0	2.03 t (9.0)
18	22.6	22.6	0.86 s	23.2	23.2	0.83 s
19	12.7	12.8	1.00 s	13.1	13.1	0.75 s
20	47.7	47.8	2.17 m	46.5	46.5	1.96 m
21	108.9	109.0	4.78 d (3.6)	104.8	104.9	4.68 d (3.6)
22	33.8	33.9	1.52 m	32.5	32.6	1.78 m/1.94 m
23	76.7	76.8	4.22 ddd (10.8, 5.4, 1.8)	77.8	77.8	4.44 ddd (9.6, 9.6, 2.4)
24	75.3	75.6	3.25 dd (9.6, 1.2)	76.7	77.0	3.26 dd (9.6, 2.4)
25	73.1	73.1	-	77.1	77.1	-
26	26.3	26.4	1.27 s	20.1	20.1	1.18 s
27	26.4	26.5	1.30 s	22.4	22.4	1.24 s
28	21.6	21.6	1.10 s	27.0	27.6	0.97 s
29	24.5	24.6	1.05 s	14.7	14.7	0.86 s
30	27.3	27.3	1.12 s	27.3	27.3	0.99 s
21-OCH ₃	55.6	55.6	3.34 s	55.0	55.0	3.35 s
25-OCH ₃				49.3	49.3	3.25 s

Measured in ^aCDCl₃, ^b150 MHz, and ^c600 MHz. [#] δ_{C} of 21 α -methylmelianodiol [16] and ^{\$} δ_{C} of 21S,23R-epoxy-3 β ,24R-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-en [13].

The strong cross-peaks observed in the HMBC spectrum (Fig. 2) between the methyl protons and their neighboring carbons were used to establish the tirucallane backbone. The location of the methoxy group at C-21 was confirmed by the HMBC cross-peak from the methoxy protons to C-21. HMBC correlations from olefinic proton H-7 [δ_{H} 5.32 (1H, dd, J = 6.6, 3.0 Hz, H-7)] to C-5 (δ_{C} 52.4), and C-9 (δ_{C} 48.4) assigned the trisubstituted double bond to C-7 (δ_{C} 118.2) and C-8 (δ_{C} 145.6). Further structural elucidation of **3** was analyzed by ^1H - ^1H COSY spectrum, which enabled the identification of four structural fragments: C-1-C-3, C-5-C-6-C-7, C-9-C-11-C-12, and C-15-C-16-C-17-C-20(C-21)-C-22-C-23-C-24 based on the sequences of correlated protons (Fig. 2). The NOESY cross-peaks for H-5/H-9, H-5/H₃-28, H-9/H₃-18, H₃-18/H-20, and H-20/H-23, indicated that H-5, H-9, H₃-18, H₃-28, H-23 were α -oriented. The observable NOESY correlations between H₃-29/H₃-19, H₃-30/H-17 suggested β -configurations for H-17, H₃-19, H₃-29 and H₃-30. The coupling constant between H-23 and H-24 (J = 9.6 Hz) in **3** indicated a *trans* relationship, which was further supported by the NOESY correlation between H-24/H-21. Furthermore, compound **3** displayed the same configuration at C-20 as compounds **1** and **2**. The comparison of the NMR data of **3** with NMR spectroscopic data in the published literature [16] supported the identification of **3**. Thus, compound **3** was established as 21 α -methylmelianodiol, previously reported from the fruits of *Poncirus trifoliata* (Rutaceae) [16] and *Melia azedarach* [17]. 21 α -Methylmelianodiol showed significant cytotoxicity against HCT116 (carcinoma) and RKO (adenocarcinoma) cell lines with IC₅₀ values of 10.16 ± 1.22 and 8.57 ± 0.80 μM , respectively [17]. The anticancer potential of **3** with a redox regulatory mechanism and also outlines the mechanistic action of **3** in inhibiting the growth, survival, and metastasis of human non-small cell lung cancer cells and influences a mechanism that results in the influence on suppression of P-gp/MDR1-associated multidrug resistance in drug-resistant human non-small cell lung cancer cells [18]. Additionally, the anti-inflammatory effect of compound **3** was evaluated based on its ability to inhibit nitric oxide (NO) production in mouse peritoneal macrophages [19]. This effect was mediated

through the suppression of interleukin-6 (IL-6) expression [20] and the inhibition of NF- κ B phosphorylation [21].

Compound **4** was obtained as a white amorphous powder with the molecular formula C₃₂H₅₄O₅, as confirmed by the sodium adduct ion peak at m/z 541.3874 [$\text{M} + \text{Na}$]⁺ (calcd. for [C₃₂H₅₄NaO₅]⁺, 541.3864). The 1D and 2D NMR spectroscopic data of **4** were highly similar to those of the compound 21 α -methylmelianodiol (**3**), indicating that both compounds shared the same tirucallane-type triterpenoid core [13],[22]. The obvious difference was the presence of the addition of a methoxy group (δ_{H} 3.25/ δ_{C} 49.3) and a ketone signal (δ_{C} 216.8) in **3** instead of an oxygenated methine signal for C-3 in **4**. In the HMBC spectrum of **4**, cross-peaks from the two *gem*-dimethyl protons at δ_{H} 0.97 and 0.86 (each 3H, s) to the oxymethine carbon resonance δ_{C} 79.2 suggested the presence of a hydroxy group at the C-3 position. Furthermore, the methoxy group was assigned to C-25 based on the HMBC correlation between OMe (δ_{H} 3.25) and C-25 (δ_{C} 77.1). The configuration of the hydroxy group was determined to be β -orientation from the multiplicity of H-3 (δ_{H} 3.23, m), in contrast to the broad singlet typically observed for an α -orientation [13],[22]. Similarly, a β -orientation of the methoxy group at C-21 was supported by the ^{13}C NMR chemical shift of C-21 in **4** (δ_{C} 104.9), which closely matched the corresponding value reported for 21*S*,23*R*-epoxy-3 β ,24*R*-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-en [13] at δ_{C} 104.8. A comprehensive analysis of the NMR spectroscopic data, together with comparison to previously published data [13],[22], led to the identification of **4** to be 21*S*,23*R*-epoxy-3 β ,24*R*-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-en.

Compound **4** was evaluated for its growth-inhibitory effects on HepG2 (liver), K562 (leukemia), SGC7901 (gastric), and HL60 (leukemia) cancer cell lines using MTT and SRB assays. The results showed that compound **4** exhibited moderate cytotoxic activity against the HL60 cell line, with an IC₅₀ value of 44.9 μM , in comparison with cisplatin (IC₅₀ = 4.2 μM) [13].

Two remaining compounds were determined to be odoratone (**5**) [23] and cabraleone (**6**) [24] by comparison of their physical and spectroscopic data to those reported in the references. Indeed, odoratone (**5**) was reported for the first time by Chan and co-workers in 1967 from *Cedrela odorata* based on its IR absorption bands and the characteristic proton signals [25].

Its ^{13}C NMR data were published for the first time by Luo et al [23]. Unfortunately, although compounds **5** and **6** have been reported in many other plants, it is worth noting that there is still little research on their biological assay.

4. Conclusions

The chemical constituents of *W. pinnata* leaves have been identified in the present study. Six compounds were isolated, including cumingianol E (**1**), 21-*O*-methyltoosendanpentol (**2**), 21 α -methylmelianodiol (**3**), 21*S*,23*R*-epoxy-3 β ,24*R*-dihydroxy-21 β ,25-dimethoxy-tirucalla-7-

en (**4**), odoratone (**5**), and cabraleone (**6**). The structures of these compounds were elucidated through NMR spectroscopic analysis and comparison with previously reported data. To the best of our knowledge, the occurrence of compounds **1–6** are reported here for the first time in this species.

Acknowledgments: This research was financially supported by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number: 104.01-2024.03.

References

1. Plants of the World Online (2025). https://powo.science.kew.org/results?f=accepted_names&q=Walsura, accessed 08/12/2025.
2. Han M. L., Zhang H., Yang S. P., Yue J. M. (2011), Walsucochinoids A and B: new rearranged limonoids from *Walsura cochinchinensis*. *Organic Letters*, 14(2), 486–489.
3. Ban N. T., Khoi N. K., Phuong V. X., Anh T. T. P., Bien L. K., Binh N. Q., Dung H. T. (2001), Checklist of plant species of Vietnam. *Agricultural Publishing House*, Vol. II, 1005–1006.
4. Son N. T. (2022), The genus *Walsura*: a rich resource of bioactive limonoids, triterpenoids, and other types of compounds. *Progress in the Chemistry of Organic Natural Products*, 118, 131–177.
5. Ji K. L., Zhang P., Hu H. B., Hua S., Liao S. G., Xu Y. K. (2014), Limonoids from the leaves and twigs of *Walsura yunnanensis*. *Journal of Natural Products*, 77(8), 1764–1769.
6. Rao M. S. A., Suresh G., Yadav P. A., Prasad K. R., Rani P. U., Rao C. V., Babu K. S. (2015), Piscidinols H–L, apotirucallane triterpenes from the leaves of *Walsura trifoliata* and their insecticidal activity. *Tetrahedron*, 71(9), 1431–1437.
7. Luo X. D., Wu D. G., Cai X. H., Kennelly E. J. (2006), New antioxidant phenolic glycosides from *Walsura yunnanensis*. *Chemistry & Biodiversity*, 3(2), 224–230.
8. Yusoff M. (2012), Chemical constituents of *Walsura pinnata* Hassk. *Master's thesis, University of Malaya, Kuala Lumpur, Malaysia*.
9. Mahdzir M. A., Shilpi J. A., Mahmud N., Ramasamy S., Awang K. (2017), Chemical constituents from *Walsura pinnata* (Meliaceae). *Natural Product Communications*, 12, 1397–1400.
10. Zakaria N., Mahdzir M. A., Yusoff M., Arshad N. M., Awang K., Nagoor N. H. (2018), Cytotoxic effects of pinnatane A extracted from *Walsura pinnata* (Meliaceae) on human liver cancer cells. *Molecules*, 23(11), e2733.
11. Azrul M. M. (2017), Phytochemical investigation of *Walsura pinnata*. *Master's thesis, University of Malaya, Kuala Lumpur, Malaysia*.
12. Kurimoto S., Kashiwada Y., Lee K. H., Takaishi Y. (2011), Triterpenes and a triterpene glucoside from *Dysoxylum cumingianum*. *Phytochemistry*, 72(17), 2205–2211.
13. Zhou F., Ma X. H., Li Z. J., Li W., Zheng W. M., Wang Z. B., Zeng X. M., Sun K. H., Zhang Y. H. (2016), Four new tirucallane triterpenoids from the fruits of *Melia azedarach* and their cytotoxic activities. *Chemistry & Biodiversity*, 13(12), 1738–1746.
14. Xie B. J., Yang S. P., Chen H. D., Yue J. M. (2007), Agladupols A–E, triterpenoids from *Aglaia duperreana*. *Journal of Natural Products*, 70(9), 1532–1535.
15. Inada A., Konishi M., Nakanishi T. (1989), Phytochemical studies on Meliaceae plants. V. Structure of a new apotirucallane-type triterpene, 21-*O*-methyltoosendanpentol from fruits of *Melia toosendan* Sieb. et Zucc. *Heterocycles*, 28, 383–387.
16. Xu G. H., Kim J. A., Kim S. Y., Ryu J. C., Kim Y. S., Jung S. H., Kim M. K., Lee S. H. (2008), Terpenoids and coumarins isolated from the fruits of *Poncirus trifoliata*. *Chemical and Pharmaceutical Bulletin*, 56(6), 839–842.
17. Song M., Chan G., Lin L. G., Li D., Zhang K., Zhang X. Q., Ye W. C., Li N., Zhang Q. W. (2022), Triterpenoids from the fruits of *Melia azedarach* L. and their cytotoxic activities. *Phytochemistry*, 201, e113280.
18. Aldonza M. B. D., Hong J. Y., Bae S. Y., Song J., Kim W. K., Oh J., Shin Y., Lee S. H., Lee S. K. (2015), Suppression of MAPK signaling and reversal of mTOR-dependent MDR1-associated multidrug resistance by 21 α -methylmelianodiol in lung cancer cells. *Plos One*, 10(7), e0133922.
19. Chen H., Ma S. G., Fang Z. F., Bai J., Yu S. S., Chen X. G., Hou Q., Yuan S. P., Chen X. (2013), Tirucallane triterpenoids from the stems of *Brucea mollis*. *Chemistry & Biodiversity*, 10(4), 695–702.
20. Lee I. J., Xu G. H., Ju J. H., Kim J. A., Kwon S. W., Lee S. H., Han S. B., Kim Y. (2008), 21-Methylmelianodiols from *Poncirus trifoliata* as inhibitors of interleukin-5 bioactivity in Pro-B cells. *Planta Medica*, 74(4), 396–400.
21. Zhou H. Y., Shin E. M., Guo L. Y., Zou L. B., Xu G. H., Lee S. H., Ze K. R., Kim E. K., Kang S. S., Kim Y. S. (2007), Anti-inflammatory activity of 21(α , β)-methylmelianodiols, novel compounds from *Poncirus trifoliata* Rafinesque. *European Journal of Pharmacology*, 572(2–3), 239–248.
22. Liu J., Yang S. P., Ni G., Gu Y. C., Yue J. M. (2012), Triterpenoids from *Aglaia odorata* var. *microphyllina*. *Journal of Asian Natural Products Research*, 14(10), 929–939.
23. Luo X. D., Wu S. H., Ma Y. B., Wu D. G. (2000), A new triterpenoid from *Azadirachta indica*. *Fitoterapia*, 71(6), 668–672.
24. Singh Y., Aalbersberg W. (1992), Dammarane triterpenoids from *Dysoxylum richii*. *Phytochemistry*, 31, 4033–4035.
25. Chan W. R., Taylor D. R., Snatzke G., Fehllhaber H. W. (1967), The tetracyclic triterpenes from *Cedrela odorata* L. *Chemical Communications*, 11, 548–550.