

ISSN 1859 - 4735



JOURNAL OF MEDICINAL MATERIALS

No. 1+2 - Vol. 31
3/2026

SPECIAL ISSUE

THE 14th SCIENTIFIC CONFERENCE OF NATIONAL INSTITUTE OF MEDICINAL MATERIALS (NIMM)
THE 65th ANNIVERSARY OF NIMM FOUNDATION (13/4/1961-13/4/2026)

NATIONAL INSTITUTE OF MEDICINAL MATERIALS

3B QUANG TRUNG - CUA NAM - HANOI - VIETNAM
TEL: (+84-24) 38247058 FAX: (+84-24) 39348740
Email: tapchiduoclieu@nimm.org.vn; tapchiduoclieu@gmail.com

EDITOR-IN-CHIEF

Assoc. Prof. PhD. Do Thi Ha

I. Subcommittee of Pharmacognosy - Traditional Medicine

• **Assoc. Prof. Dr. Sci. Nguyen Minh Khoi - Head of Subcommittee**

- Assoc. Prof. PhD. Nguyen Thuong Dong - Member
- Prof. PhD. Nguyen Minh Duc - Member
- Prof. PhD. Tran Cong Luan - Member
- Prof. PhD. Won Keun Oh - Member
- Assoc. Prof. PhD. Tran Hung - Member
- Assoc. Prof. PhD. Nguyen Thi Bich Thu - Member
- Assoc. Prof. PhD. Phuong Thien Thuong - Member
- Assoc. Prof. PhD. Phan Minh Giang - Member
- Assoc. Prof. PhD. Le Nguyen Thanh - Member
- Assoc. Prof. PhD. Le Viet Dung - Member
- Assoc. Prof. PhD. Nguyen Quoc Huy - Member
- Assoc. Prof. PhD. Nguyen Thu Hang - Member
- PhD. Tran Minh Ngoc - Member
- PhD. Nguyen Van Tai - Member
- Assoc. Prof. PhD. Le Dinh Chi - Member
- PhD. Nguyen Tuan Hiep - Member.

II. Subcommittee of Pharmacology - Clinical Pharmacy

• **Assoc. Prof. PhD. Pham Thi Nguyet Hang - Head of Subcommittee**

- Assoc. Prof. Dr.Sci. Do Trung Dam - Member
- Assoc. Prof. PhD. Nguyen Thi Thu Huong - Member
- Assoc. PhD. William R. Folk - Member
- Assoc. Prof. PhD. Pham Thi Van Anh - Member
- Assoc. Prof. PhD. Nguyen Thuy Duong - Member
- Assoc. Prof. PhD. Le Van Minh - Member
- PhD. Tran Thi Hien - Member.

III. Subcommittee of Agriculture - Biology

• **PhD. Nguyen Van Khiem - Head of Subcommittee**

- PhD. Phan Thuy Hien - Member
- Assoc. Prof. PhD. Nguyen Van Tap - Member
- Assoc. Prof. PhD. Pham Thanh Huyen - Member
- Prof. PhD. Tran The Bach - Member.

IV. Subcommittee of Secretariat Administration

• **Assoc. Prof. PhD. Le Nguyen Thanh - Head of Subcommittee**

- PhD. Le Thi Xoan - Academic Secretary
- BSc. Nguyen Thi Minh Loc - Administrative Secretary.

CONTENTS VOL.31(1+2), 2026 (SPECIAL ISSUE)
(Pages 1 - 160)

Page	SCIENTIFIC RESEARCH
3	Foreword: Special Issue Commemorating the 65th Anniversary of the National Institute of Medicinal Materials and Welcoming the 1st International Conference V.IMPACT-2026 - Assoc. Prof. Do Thi Ha, PhD
	REVIEW
4	Results of Scientific Activities of the National Institute of Medicinal Materials (2016–2025) and Development Orientation Toward 2030 - Tran Minh Ngoc, Do Thi Ha, Pham Tuan Anh
	SCIENTIFIC RESEARCH
10	Triterpenoid Constituents from the Leaves of <i>Walsura pinnata</i> Hassk - Bui Huu Tai, Nguyen Viet Dung, Nguyen Huy Hoang, Ngo Anh Bang, Duong Thi Dung, Duong Thi Hai Yen, Phan Thi Thanh Huong, Dan Thuy Hang, Pham Hai Yen, Phan Van Kiem, Nguyen Phuong Thao
17	Chemical Constituents from the Propolis of the Stingless Bee <i>Lepidotrigona terminata</i> (Smith, 1878) Collected in Gia Lai - Diep Thi Lan Phuong, Nguyen Thu Thao Van, Vo Thi Thanh Tuyen, Nguyen Thi Nghia, Pham Ngoc Thach, Le Thi Cam Nhung, Le Duy Thanh, Vo Manh Tien, Tran Minh Ngoc, Le Nguyen Thanh
23	Glycoside Constituents from <i>Abroma augustum</i> and their <i>in vitro</i> Antiproliferative Activity - Ninh Thi Ngoc, Pham Thi Cham, Le Thi Vien, Tran Thi Hong Hanh, Do Thi Thao, Nguyen Xuan Cuong, Nguyen Hoai Nam
29	Chemical Profiling and Anti-Inflammatory Effects of <i>Physalis angulata</i> in Lipopolysaccharide-Stimulated Bone Marrow–Derived Dendritic Cells - Dinh Thi Huong, Le Ba Vinh, Nguyen Van Chinh, Nguyen Cao Cuong
36	Phenolic Constituents from the Ethyl Ecetate Extract of <i>Camellia flava</i> Flowers (Theaceae) Collected in Phu Tho Province - Do Thi Thuy, Nguyen Duong Hong Anh, Tran The Bach, Do Van Hieu, Do Duc Nhuan, Than Kieu My, Tran Minh Ngoc, Nguyen Thanh Cong, Le Nguyen Thanh
42	Lignans and Other Compounds from the Leaves of <i>Triadica rotundifolia</i> with Inhibitory Activity on Nitric Oxide Production - Ninh Khac Ban, Tran My Linh, Ninh Khac Thanh Tung, Bui Van Thanh, Pham Hai Yen, Bui Huu Tai, Phan Van Kiem, Nguyen Phuong Thao
49	Chemical Constituents of <i>Ganoderma multipileum</i> Ding Hou, Ganodermataceae - Nguyen Thi Duyen, Tran Thi Tuyet, Phan Quynh Trang, Nguyen Trung Hieu, Nguyen Duc Hieu, Nguyen Phuong Dai Nguyen
53	Chemical Composition, Antimicrobial and Cytotoxic Activities of the Leaf Essential Oil of <i>Knema globularia</i> (Lamk.) Warb. from Thanh Hoa, Vietnam - Luong Le Uyen Trang, Pham Thi Huong Ly, Nguyen Thi Hoa Hien, Nguyen Thanh Huyen, Nguyen Thi Hien, Nguyen Thi Thu, Khong Thi Hoa

60	Azaphilone Derivatives Isolated from Endophytic Fungus <i>Penicillium sclerotiorum</i> - Ly Tu Loan, Vo Van Leo, Ma Chi Thanh
67	Chemical Constituents from the Aerial Parts of <i>Barleria lupulina</i> (Acanthaceae) - Ngo Thi Ngoc Yen, Do Hoang Anh, Nguyen Thi Thu, Nguyen Thi Ha Ly, Le Nguyen Thanh, Nguyen Quynh Nga, Dao Viet Quoc, Nguyen Thu Hang, Nguyen Hoang Minh, Nguyen Minh Khoi, Le Tran Nguyen Vu
74	Determination of Essential Oil Content, Composition, and Total Polyphenol Content of <i>Rhizoma homalomenae occulta</i> Collected in Vietnam - Nguyen Thi Ha Ly, Pham Thi Hien, Ngo Thi Thanh Huong, Hoang Dieu Linh, Tran Huyen Trang, Nguyen Minh Khoi
80	Establishment of an In-House Standard for <i>Prunellae spica</i> - Tran Huyen Trang, Pham Thi Hien, Nguyen Thi Ha Ly, Phan Van Truong, Vu Thi Diep, Nguyen Tra My, Nguyen Thi Hang, Nguyen Duong Hong Anh, Do Thi Ha
88	HPTLC Scanner Method for the Simultaneous Determination of Lycopene and β-Carotene in Gac Seed Aril - Pham Thi Hien, Pham Thi Khuyen, Nguyen Thi Hang, Nguyen Thi Lan Hoa, Ta Thi Thao, Do Thi Ha, Nguyen Thi Ha Ly
96	Optimization of the Extraction Process from Yellow Camellia Flowers (<i>Camellia cucphuongensis</i> Ninh & Rosmann) Collected in Phu Tho Province Using Response Surface Methodology - Nguyen Hoang Viet, Nguyen Dinh Quan, Do Duc Nhuan, Nguyen Duong Hong Anh, Tran Thi Mai Anh, Doan Van Nghia, Vu Xuan Hoa, Dang Mai Thien Ha, Nguyen Thanh Cong
105	Antioxidant, Antihyperglycemic, and Renal Protective Effects of <i>Hibiscus tiliaceus</i> L. Leaf Extract on Streptozotocin-Induced Hyperglycemia - Nguyen Thi Thu Huong, Nhieu Chinh Hoang, Nguyen Hoang Minh
113	Acute Oral Toxicity and Laxative Effects of <i>Pharbitis nil</i> (L.) Choisy Seed Extract - Nguyen Thi Kim Oanh, Tran Vien Thong, Truong Huy Hieu, Ly Hai Trieu, Pham Thi Nguyet Hang, Nguyen Thi Vinh Hue, Do Thi Ha
123	Evaluation of Memory-Enhancing Effects of <i>Morinda officinalis</i> How. Root Extracts in a Transgenic <i>hAPP Drosophila</i> Model of Alzheimer's Disease - Nguyen Van Hiep, Pham Thi Nguyet Hang, Trinh Dinh Luc, Leu Khanh Duy, Pham Anh Tung, Nguyen Thi Phuong, Nguyen Van Tai
131	Anti-Inflammatory Effects of the Bark of <i>Phellodendron amurense</i> Rupr. in Experimental Models - Do Thi Hang, Le Thi Kim Van, Do Van Dung, Bui Thanh Tung
138	Neuroprotective Effects of Tianma Gouteng Decoction in a Middle Cerebral Artery Occlusion Model of Ischemic Stroke - Le Thi Xoan, Trinh Lan Phuong, Nguyen Van Hiep, Pham Thi Nguyet Hang, Nguyen Van Tai, Nguyen Thi Thu Trang, Nguyen Thi Thanh Loan, Nguyen Hung Son
147	Investigation of the Variation of Gymnemagenin Content in <i>Gymnema R.Br</i> Under the Influence of Certain Strains of Bacteria - Tong Thi Thanh Vuong, Tran Thi Minh Thuy, Le Thi Kim Van
154	Comparative Transcriptome Profiling of <i>Momordica cochinchinensis</i> (Lour.) Spreng. Aril Using Oxford Nanopore and Illumina Sequencing Technologies - Tran Duc Trung, Nguyen Hoang, Le Thi Tu Linh, Nguyen Thi Thu Hang, Pham Thanh Huyen, Truong Nhat My, Nguyen Thi Lan Hoa

**CHEMICAL CONSTITUENTS FROM THE AERIAL PARTS
OF *BARLERIA LUPULINA* (ACANTHACEAE)**

**Ngo Thi Ngoc Yen^{1,2}, Do Hoang Anh¹, Nguyen Thi Thu¹, Nguyen Thi Ha Ly¹, Le Nguyen Thanh¹,
Nguyen Quynh Nga¹, Dao Viet Quoc¹, Nguyen Thu Hang³, Nguyen Hoang Minh³,
Nguyen Minh Khoi^{1,*}, Le Tran Nguyen Vu^{3,*}**

¹National Institute of Medicinal Materials, Hanoi 11018, Vietnam;

²Faculty of Medicine and Pharmacy, Tay Nguyen University, Vietnam;

³Research Center of Ginseng and Medicinal Materials (CGMM), National Institute of Medicinal Materials, Ho Chi Minh City 71016, Vietnam

*Corresponding author: khoi_nguyenminh@yahoo.co.uk and nguyenvu19hc@gmail.com.

Received February 10th, 2026

Accepted March 13th, 2026

Summary

Phytochemical studies of the *n*-hexane and ethyl acetate fractions from aerial parts of *Barleria lupulina* L. (Acanthaceae) collected in Cu Chi, Ho Chi Minh city, Vietnam, led to the isolation and identification of six known compounds, including (+)-lyoniresinol (1), (+)-lyoniresinol 9'-*O*- β -D-glucopyranoside (2), 2-methyl-1,3,6-trihydroxy-9,10-anthraquinone 3-*O*- β -D-glucopyranoside (3), (3*S*)-l-octene-3-ol 3-*O*- β -D-glucopyranoside (4), uracil (5) and stigmasterol (6). The structures of the isolated compounds were determined on the basis of detailed spectroscopic (MS and NMR) analysis and comparison with reported literature. Compounds 1, 3, 4, and 5 were found for the first time from the *Barleria* genus.

Keywords: *Barleria lupulina*; Lignan; Anthraquinone; Sterol.

1. Introduction

Genus *Barleria* (Acanthaceae), which comprises over 300 species of herbs and shrubs, has been traditionally utilized across various countries for treating a wide range of ailments [1]. *Barleria lupulina* L., commonly known as hop-headed barleria, is one species recognized for its decorative and medicinal value. Traditionally, different parts of the plant have been employed in Asian countries, such as India, Thailand, and Indonesia, for the treatment of diabetes, rheumatoid arthritis, inflammation, skin infections, and mental strain [2]. In Vietnam, the leaves of *B. lupulina* have been applied in case of snakebites and fever [3]. Recent phytochemical studies have revealed that *B. lupulina* is a rich source of diverse bioactive compounds, predominantly iridoid glycosides, phenylethanoid glycosides, phenylpropanoid glycosides, and phenolic compounds [2],[4],[5],[6]. These isolated constituents are associated with a broad spectrum of pharmacological activities, including antibacterial, anti-inflammatory, anti-arthritis, anti-diabetic, antioxidant, anti-osteoporotic, anti-ulcer, and neuroprotective effects [2],[7],[8],[9].

Despite promising traditional uses and preliminary scientific validation, the chemical investigations of *B. lupulina* in Vietnam are still

limited [5],[6],[7]. Preliminary biological screening results showed that the *n*-hexane and ethyl acetate extracts from the *B. lupulina* sample collected in Ho Chi Minh city displayed potent anti-inflammatory activity, inhibiting 72-87% of NO-production in LPS-induced RAW264.7 cells at the concentration of 50 μ g/mL. Hence, our research paper aims to provide an isolation and identification of chemical constituents from *n*-hexane and ethyl acetate extracts, including (+)-lyoniresinol (1), (+)-lyoniresinol 9'-*O*- β -D-glucopyranoside (2), 2-methyl-1,3,6-trihydroxy-9,10-anthraquinone 3-*O*- β -D-glucopyranoside (3), 3-*O*- β -D-glucopyranosyl (3*S*)-l-octene-3-ol (4), uracil (5), and stigmasterol (6).

2. Materials and methods

2.1. Materials

The aerial parts of *Barleria lupulina* L. (Acanthaceae) were collected in Cu Chi, Ho Chi Minh City, in 2024. The plant species was identified by MSc Nguyen Quynh Nga and BS. Dao Viet Quoc, Center of Medicinal Material Resources (CMMR), National Institute of Medicinal Materials. The voucher specimen (NIMM-20133) was deposited at the National Institute of Medicinal Materials.

2.2. General experimental procedures

ESI-MS spectra were acquired on an Agilent

1260 series single quadrupole LC/MS system. NMR spectra (^1H , ^{13}C , HSQC, HMBC, and NOESY) were obtained on Bruker AVANCE III HD 500 MHz or Bruker AVANCE NEO 600 MHz spectrometers with tetramethylsilane as the internal standard. Column chromatography (CC) was performed on normal phase *silica gel* 60 (230–400 mesh, Merck), reversed phase (RP-18) resin (YMC), and Sephadex® LH20. Preparative HPLC was conducted by Shimadzu HPLC (LC-20AP), SunFire® C18 OBD™ Prep Column (100 Å, 10 μm , 10 $\times$ 250 mm). Thin-layer chromatography has been performed on *silica gel* 60 GF₂₅₄ and RP-18 G F₂₅₄S. Compounds were visualized by spraying with 10% sulfuric acid in EtOH and heating.

2.3. Extraction and isolation

The dried aerial parts (28 kg) were macerated and extracted with 70% EtOH (3 times $\times$ 220 L for 24 h) at room temperature. The combined extracts were concentrated under reduced pressure to yield the EtOH extract (4.3 kg). The EtOH extract (4 kg) was suspended in 4 L of distilled water and then successively extracted with *n*-hexane, dichloromethane (DCM), and EtOAc (each 3 $\times$ 4 L). The organic solvents were then removed *in vacuo* to afford *n*-hexane (95 g), DCM (265 g), and EtOAc (300 g) extracts, respectively.

The *n*-hexane extract (90 g) was chromatographed on a *silica gel* CC, eluted with a solvent gradient of *n*-hexane–acetone (20:1 to 1:1, v/v), DCM–MeOH (20:1 $\rightarrow$ 2:1, v/v), and MeOH to give 11 fractions (H1–H11). Fraction H1 (2.8 g) was chromatographed by *silica gel* CC, eluted with a gradient solvent of *n*-hexane–acetone (15:1 $\rightarrow$ 1:1, v/v) to afford 3 fractions H1.1–H1.3. Compound **6** (50 mg) was obtained by recrystallization in MeOH from fraction H1.3.

The EtOAc extract (240 g) was applied to *silica gel* CC, eluted with a solvent gradient of *n*-hexane–acetone (20:1 to 1:1, v/v), DCM–MeOH (20:1 $\rightarrow$ 2:1, v/v), and MeOH to give 10 fractions (E1–E10). Fraction E6 (3.0 g) was chromatographed by Sephadex® LH20 CC (MeOH–H₂O 1:1, v/v), yielding 5 fractions E6.1–E6.5. Fraction E6.2 (1.8 g) was separated by *silica gel* CC eluted with a solvent gradient of

DCM–MeOH (20:1 $\rightarrow$ 2:1, v/v) to afford ten fractions E6.2.1–E6.2.10. Fraction E6.2.4 (55 mg) was then purified by reversed-phase *silica gel* CC (MeOH–H₂O 1:1, v/v) to afford **1** (3.0 mg). Fraction 6.2.7 (89 mg) was subjected to a reversed-phase *silica gel* CC (MeOH–H₂O 2:3, v/v), yielding 2 sub-fractions E6.2.7.1 and E6.2.7.2. Fraction 6.2.7.1 (32 mg) was further purified by *silica gel* CC (EtOAc–MeOH 35:1, v/v) to give **5** (4.1 mg). Fraction E6.2.8 (138 mg) was chromatographed by reversed-phase *silica gel* CC (MeOH–H₂O 1:1, v/v) to afford **3** (9.0 mg). Fraction E8 (23.4 g) was subjected to *silica gel* CC, eluted with gradient solvents of DCM–MeOH (10:1 - 100% MeOH, v/v), yielding 12 fractions E8.1–E8.12. Fraction E8.9 (6.2 g) was purified by reversed-phase *silica gel* CC (MeOH–H₂O 1:2, v/v) to give 12 sub-fractions E8.9.1–E8.9.12. Fraction E8.9.2 (3.2 g) was separated by reversed-phase *silica gel* CC (*n*-hexane–acetone v/v, 19:1), followed by Sephadex LH-20 to afford 3 fractions (E8.9.2.1–E8.9.2.3). Fraction 8.9.2.1 (170 mg) was purified by preparative HPLC (acetonitrile–H₂O 13:87 (v/v), flow rate: 2.5 mL/min, injection volume 65 μL , detection wavelength 210 nm) to yield compound **2** (2.6 mg, t_{R} = 45 min). Fraction 8.9.2.3 (55 mg) was purified by preparative HPLC (MeOH–H₂O 43:57 (v/v), flow rate: 2.5 mL/min, injection volume 49 μL , detection wavelength 254 nm) to give **4** (14 mg, t_{R} = 15 min).

(+)-Lyoniresinol (1): White solid; ESI-MS: m/z 419 $[\text{M}-\text{H}]^-$; $[\alpha]_{\text{D}}^{20} = +17.9$ (MeOH, $c = 0.56$). ^1H -NMR (600 MHz, CD₃OD) and ^{13}C -NMR (125 MHz, CD₃OD): See **Table 1**.

(+)-Lyoniresinol 9'-O- β -D-glucopyranoside (2): White solid; $[\alpha]_{\text{D}}^{20} = +60.1$ (MeOH, $c = 0.33$). ESI-MS m/z 581 $[\text{M}-\text{H}]^-$. ^1H -NMR (600 MHz, CD₃OD) and ^{13}C -NMR (125 MHz, CD₃OD): See **Table 1**.

2-Methyl-1,3,6-trihydroxy-9,10-anthraquinone 3-O- β -D-glucopyranoside (3): Yellow solid, ESI-MS: m/z 291 $[\text{M}+\text{H}]^+$. ^1H -NMR (600 MHz, CD₃OD) and ^{13}C -NMR (125 MHz, CD₃OD): See **Table 2**.

3-O- β -D-Glucopyranosyl (3S)-l-octene-3-ol (4): Yellow solid, $[\alpha]_{\text{D}}^{20} = -7.9$ (MeOH, $c = 1.27$). ^1H -NMR (600 MHz, CD₃OD) and ^{13}C -NMR (125 MHz, CD₃OD): See **Table 2**.

Uracil (5): Yellow solid. ESI-MS m/z 111.2 $[M-H]^-$. 1H -NMR (500 MHz, CD_3OD): δ_H 5.61 (1H, d, $J = 7.5$ Hz, H-5), 7.38 (1H, d, $J = 7.5$ Hz, H-6). ^{13}C -NMR (125 MHz, CD_3OD): δ_C 151.5 (C-2), 167.3 (C-4), 101.7 (C-5), 143.5 (C-6).

Stigmasterol (6): White solid, ESI-MS m/z 413 $[M+H]^+$. 1H -NMR (600 MHz, $CDCl_3$): δ_H 3.52 (1H, m, H-3), 5.34 (1H, m, H-6), 0.70 (3H, s, H-18), 1.01 (3H, s, H-19), 1.03 (3H, d, $J = 6.6$ Hz, H-21), 5.15 (1H, dd, $J = 15.0$; 8.4 Hz, H-22), 5.02 (1H, dd, $J = 15.0$; 8.4 Hz, H-23), 0.84 (3H, d,

$J = 6.6$ Hz, H-26), 0.81 (3H, d, $J = 6.6$ Hz, H-27), 0.80 (3H, t, $J = 7.2$ Hz, H-29). ^{13}C -NMR (125 MHz, $CDCl_3$): δ_C 37.3 (C-1), 31.7 (C-2), 71.8 (C-3), 42.3 (C-4), 140.8 (C-5), 121.7 (C-6), 31.9 (C-7), 31.9 (C-8), 50.2 (C-9), 36.5 (C-10), 21.1 (C-11), 39.7 (C-12), 42.3 (C-13), 56.9 (C-14), 24.4 (C-15), 28.9 (C-16), 56.0 (C-17), 12.1 (C-18), 19.4 (C-19), 40.5 (C-20), 21.1 (C-21), 138.3 (C-22), 129.3 (C-23), 51.3 (C-24), 31.9 (C-25), 21.2 (C-26), 19.0 (C-27), 25.4 (C-28), 12.2 (C-29).

3. Results and discussion




Fig. 1. Chemical structures of compounds 1–6 from *B. lupulina*

Compound **1** was isolated as a white solid. The 1H -, ^{13}C -NMR and HSQC spectra of **1** exhibited signals of 3 aromatic protons at δ_H 6.61 (1H, s, H-2)/ δ_C 107.8 (C-2), δ_H 6.40 (2H, s, H-2' and H-6')/ δ_C 106.98 (C-2); 2 oxymethylene groups at δ_H 3.61–3.51 (2H, H-9)/ δ_C 66.8 (C-9) and δ_H 3.51 (2H, d, $J = 5.4$ Hz, H-9')/ δ_C 64.2 (C-9'), 4 methoxy groups at δ_H 3.40 (3H, s, 3-OMe)/ δ_C 56.6; δ_H 3.88 (3H, s, 5-OMe)/ δ_C 60.2 and 3.76 (6H, s, 3'-OMe, 5'-OMe)/ δ_C 56.8; one methylene group at δ_H 2.72–2.59 (2H, H-7)/ δ_C 33.6 (C-7), 3 methine groups at δ_H 1.64 (1H, m, H-8)/ δ_C 40.9 (C-8), δ_H 4.33 (1H, d, $J = 5.4$ Hz, H-7')/ δ_C 42.3 (C-7') and δ_H 1.99 (1H, m, H-8')/ δ_C 49.9 (C-8'). The negative ESI-MS spectrum of **1** showed a deprotonated molecular ion peak at m/z 419 $[M-H]^-$, which gave evidence of its molecular formula as $C_{22}H_{28}O_8$ ($M = 420$). The

NMR and MS spectra suggested that **1** was an aryltetralin lignan. The aromatic proton at C-2 was confirmed by the HMBC correlations of H-2 (δ_H 6.61) to C-7 (δ_C 33.6) and of H-7 (δ_H 2.72 and 2.59) to C-2 (δ_C 107.8). The symmetrical 3',5'-dimethoxy-4'-hydroxy phenyl group substituted at C-7' was confirmed by the HMBC correlations of H-2' and H-6' (δ_H 6.40) to C-7' (δ_C 42.3) and C-4' (δ_C 134.6), of methoxy protons (δ_H 3.76) to C-3' and C-5' (δ_C 149.0). In addition, the methoxy groups at C-3 and C-5 were assigned by HMBC cross-peaks of methoxy protons at δ_H 3.40 and 3.88 to C-3 (δ_C 56.6) and C-5 (δ_C 60.2), respectively. In addition, HMBC cross-peaks of H-7 (δ_H 2.72 and 2.51) to C-8' (δ_C 49.9) and C-9 (δ_C 66.8), of H-7' (δ_H 4.33) to C-9' (δ_C 64.2) and C-8 (δ_C 40.9) confirmed the aryltetralin skeleton (Fig. 2). The relative configuration of **1** was

assigned based on the NOESY spectrum. The cross-peaks of H-8' (δ_{H} 1.99) with H-2' (δ_{H} 6.40) and H-9 (δ_{H} 3.51 and 3.61); of H-7' (δ_{H} 4.33) with H-8 (δ_{H} 1.64) and H-9' (δ_{H} 3.51) suggested H-7', H₂-9' and H-8 are on the same side, while the

phenyl ring, H-8' and H₂-9 are on the other side. Based on spectral analysis and the specific optical rotation value of +17.9 (ref [10]: $[\alpha]_{\text{D}}^{25} = +65.1$ (MeOH, c 0.10)), **1** was identified as (+)-lyoniresinol [11].




Fig. 2. Key HMBC (→) correlations of compounds **1**, **3**, and **4**

Compound **2** was isolated as a white solid. The molecular formula of **2** was determined as C₂₈H₃₈O₁₃ based on a deprotonated molecular ion peak at m/z 581 [M-H]⁻ in the ESI-MS spectrum and NMR data. The ¹H-, ¹³C-NMR and HSQC spectra of **2** revealed similar signals as those of **1** including 3 aromatic protons (δ_{H} 6.58 (1H, s, H-2)/ δ_{C} 107.9 (C-2), δ_{H} 6.43 (2H, s, H-2' and H-6')/ δ_{C} 107.0 (C-2', C-6')); 2 oxymethylene groups (δ_{H} 3.64-3.53 (2H, H-9)/ δ_{C} 66.3 (C-9) and δ_{H} 3.88-3.45 (2H, H-9')/ δ_{C} 71.5 (C-9')), one methylene (δ_{H} 2.71-2.61 (2H, H-7)/ δ_{C} 33.8 (C-7)) and 3 methine groups (δ_{H} 1.70 (1H, m, H-8)/ δ_{C} 40.6 (C-8), δ_{H} 4.41 (1H, d, J = 6.6 Hz, H-7')/ δ_{C}

42.8 (C-7') and δ_{H} 2.08 (1H, m, H-8')/ δ_{C} 46.7 (C-8')). The NMR spectra of **2** showed additional signals of a β -glucopyranosyl moiety at δ_{H} 4.28 (1H, d, J = 7.8 Hz, H-1''), δ_{H} 3.24-3.83 (5H, H-2''-H-6'')/ δ_{C} 104.8 (C-1''), 75.2 (C-2''), 78.0 (C-3''), 71.7 (C-4''), 78.3 (C-5''), 62.9 (C-6''). The HMBC correlations of H-1'' (δ_{H} 4.28) with C-9' (δ_{C} 71.5) indicated that the β -D-glucopyranoside moiety is linked at C-9'. Compound **2** was assigned as (+)-lyoniresinol 9'-O- β -D-glucopyranoside by comparison of specific optical rotation (+60.1, Ref. [12] $[\alpha]_{\text{D}}^{20} = +47$ (MeOH, c = 0.8)) and NMR data with previous literature [11].

Table 1. ¹H- and ¹³C-NMR spectroscopic data of **1**–**2**

Pos.	1	Ref [11]		2	Ref [11]	
	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,c}}$
1		130.2	130.2		130.2	130.2
2	6.61 (1H, s)	107.8	107.8	6.58 (1H, s)	107.9	107.9
3		148.7	148.7		148.7	148.7
4		138.9	138.9		138.9	138.9
5		147.7	147.7		147.6	147.6
6		126.3	126.3		126.4	126.4

Pos.	1		Ref [11]	2		Ref [11]
	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,c}}$
7	2.72 (1H, dd, 15.0, 4.8) 2.59 (1H, dd, 15.0, 11.4)	33.6	33.6	2.71 (1H, dd, 15.0, 4.8) 2.61 (1H, dd, 15.0, 12.0)	33.8	33.8
8	1.64 (1H, m)	40.9	40.9	1.70 (1H, m)	40.6	40.6
9	3.61 (1H, dd, 9.0, 5.4) 3.51 (1H, m)	66.8	66.8	3.64 (1H, m) 3.53 (1H, dd, 10.8, 6.6)	66.3	66.2
3-OCH ₃	3.88 (3H, s)	56.6	56.6	3.86 (3H, s)	56.6	56.6
5-OCH ₃	3.40 (3H, s)	60.2	60.2	3.35 (3H, s)	60.2	60.2
1'	-	139.3	139.3		139.3	139.3
2', 6'	6.40 (2H, s)	106.9	106.9	6.43 (2H, s)	107.0	107.0
3', 5'	-	149.0	149.0		149.0	149.0
4'	-	134.6	134.5		134.5	134.5
7'	4.33 (1H, d, 5.4)	42.3	42.3	4.41 (1H, d, 6.6)	42.8	42.8
8'	1.99 (1H, dq, 9.0, 5.4)	49.9	49.9	2.08 (1H, m)	46.7	46.7
9'	3.51 (2H, d, 5.4)	64.2	64.2	3.88 (1H, dd, 9.6, 5.4). 3.45 (1H, dd, 9.6, 4.2)	71.5	71.5
3',5'-OCH ₃	3.76 (6H, s)	56.8	56.8	3.75 (6H, s)	56.9	56.9
1'				4.28 (1H, d, 7.8)	104.8	104.8
2'				3.24 (1H, m)	75.2	75.2
3'				3.24 (1H, m)	78.0	77.9
4'				3.28 (1H, m)	71.7	71.7
5'				3.36 (1H, m)	78.3	78.2
6'				3.83 (1H, dd, 12.0, 2.4) 3.64 (1H, m)	62.9	62.8

^a: CD₃OD, ^b 600 MHz, ^c 125 Mz.

Compound **3** was isolated as a pale yellow solid. The negative ESI-MS spectrum showed the peak m/z 323 [M-H]⁻, suggesting the molecular formula C₂₁H₂₀O₁₀. The ¹H-NMR spectrum of **3** displayed 4 aromatic signals including a singlet at δ_{H} 7.41 (1H, s) and 3 protons of an ABX system at δ_{H} 7.39 (1H, d, J = 2.4 Hz, H-5), 7.03 (1H, dd, J = 2.4, 8.4 Hz, H-7), and 8.03 (1H, d, J = 8.4 Hz, H-8); a methyl singlet at δ_{H} 2.18 (3H, s, 2-Me); and signals of a sugar at δ_{H} 5.07 (1H, d, J = 2.4 Hz, H-1'), 3.37-3.43 (4H, m, H-2', H-3', H-4', H-5'), and 3.82 (1H, dd, J = 12.0; 2.4 Hz, H-6a), 3.67 (1H, dd, J = 12.0, 6.0 Hz, H-6b). The ¹³C NMR and HSQC spectra showed signals for 21 carbons, including 2 ketone groups at δ_{C} 188.1

(C-9) and 183.8 (C-10); 12 aromatic carbons, a methyl carbon at δ_{C} 8.6 and carbon signals of a β -D-glucopyranosyl part at δ_{C} 101.6 (C-1'), 74.8 (C-2'), 78.1 (C-3'), 71.1 (C-4'), 78.3 (C-5'), and 62.3 (C-6'). The NMR and MS data suggested that **3** was an anthraquinone glucoside. HMBC correlations of methyl protons to C-1 (δ_{C} 163.2), C-2 (δ_{C} 122.5), and C-3 (δ_{C} 162.1); of H-4 (δ_{H} 7.41) to C-10 (δ_{C} 183.8) and C-2; of H-8 (δ_{H} 8.03) to C-9 (δ_{C} 188.1) allowed the assignment of substitutions at the anthraquinone skeleton. Comparing the NMR data with reference [13], compound **3** was determined as 2-methyl-1,3,6-trihydroxy-9,10-anthraquinone 3-*O*- β -D-glucopyranoside.

Table 2. ¹³C-NMR spectroscopic data of **3–4**

Pos.	3		Ref [13]	4		Ref [14]
	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,d}}$	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,c}}$
1	-	163.2	163.4	5.19 (1H, d, 16.5) 5.08 (1H, d, 11.0)	141.0	141.0
2	-	122.5	120.7	5.87 (1H, ddd, 16.5, 11.0, 7.0)	116.0	116.0
3	-	162.1	160.7	4.12 (1H, q, 7.0)	82.8	82.8
4	7.41 (1H, s)	106.9	105.6	1.68 (1H, m) 1.52 (1H, m)	33.0	33.0
5	7.39 (1H, d, 2.4)	114.1	112.5	1.40–1.29 (6H, m)	25.6	25.6
6	-	165.8	161.2		35.6	35.6
7	7.03 (1H, dd, 8.4, 2.4)	122.5	121.3		23.6	23.6
8	8.03 (1H, d, 8.4)	130.7	129.5	0.90 (3H, t, 7.0)	14.4	14.4

Pos.	3		Ref [13]	4		Ref [14]
	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	$\delta_{\text{C}}^{\text{a,d}}$	$\delta_{\text{H}}^{\text{a,b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,c}}$	δ_{C}
9	-	188.1	186.3			
10	-	183.8	181.5			
11	-	137.0	135.2			
12	-	126.0	124.4			
13	-	112.0	110.5			
14	-	133.7	131.9			
2-Me	2.18 (3H, s)	8.6	8.3			
1'	5.07 (1H, d, 7.2)	101.6	100.3	4.32 (1H, d, 7.5)	103.2	103.2
2'		74.8	73.1	3.19 (1H, m)	75.3	75.3
3'		78.1	76.2	3.32 (1H, m)	78.2	78.2
4'	3.37–3.47 (4H, m)	71.1	69.3	3.29 (1H, m)	71.7	71.6
5'		78.3	72.2	3.19 (1H, m)	77.8	77.8
6'	3.82 (1H, dd, 12.0, 2.4) 3.67 (1H, dd, 12.0, 6.0)	62.3	60.4	3.80 (1H, dd, 12.0, 2.4) 3.65 (1H, dd, 12.0, 6.0)	62.8	62.8

^a: CDCl_3 , ^b: 600 MHz, ^c: 125 MHz, ^d: 100 MHz.

Compound **4** was isolated as a yellow solid. The ^1H -NMR spectrum of **4** shows signals of allyl group at δ_{H} 5.19 (1H, d, J = 16.5 Hz, H-1a), 5.08 (1H, d, J = 11.0 Hz, H-1b), 5.87 (1H, ddd, J = 16.5, 11.0, 7.0 Hz, H-2), an oxymethine group at δ_{H} 4.12 (1H, q, J = 7.0 Hz, H-3), 4 methylene groups at δ_{H} 1.68–1.52 (2H, m, H-4), 1.40–1.29 (6H, m, H-5, H-6, H-7), and a methyl group at δ_{H} 0.90 (3H, t, J = 7.0 Hz). In addition, a β -glucose moiety was observed with signals at δ_{H} 4.32 (1H, d, J = 7.5 Hz, H-1), 3.19–3.32 (4H, m, H-2, H-3, H-4, H-5), 3.80 (1H, dd, J = 12.0, 2.4 Hz, H-6a) and 3.65 (1H, dd, J = 12.0, 6.0 Hz, H-6b). The HMBC correlations of H-1' (δ_{H} 4.32) and H-1 (δ_{H} 5.19 and 5.08) to C-3 (δ_{C} 82.8) indicated the sugar linked to the carbon chain at C-3. Compound **4** was identified as (-)-3-*O*- β -D-glucopyranosyl (3*S*)-1-octen-3-ol by comparison of specific optical rotation ($[\alpha]_{\text{D}}^{20}$ = -37.2 (MeOH, c = 0.17) and NMR data with previous literature [14].

Finally, compounds **5** and **6** were determined as uracil and stigmasterol, respectively, based on the comparison of NMR data with those in the previous literature [15], [16].

(+)-Lyoniresinol 9'-*O*- β -glucoside **2** has already been identified from *B. lupulina* [1], but its aglycone (**1**) was found in the genus *Barleria* for the first time. (+)-Lyoniresinol (**1**) was identified in various plants and displayed antioxidant, anti-inflammatory, and cytoprotective activities [17], while (+)-lyoniresinol 9'-*O*- β -glucoside (**2**) exhibited antimicrobial and antidiabetic effects [18]. In the assay, (+)-lyoniresinol showed weak

NO-production inhibitory activity with an IC_{50} value of 238 μM [19]. Compound **3** was only reported from *Rubia cordifolia* (Rubiaceae family); it was first identified from the *Barleria* genus. Compound **4** might be a hydrolyzed product from 1-octen-3-yl β -primeveroside, which was found in this plant [1]. Uridine (**5**) is an essential nucleoside that plays a pivotal role in various biological processes, including macromolecule synthesis, circadian rhythms, inflammatory response, and antioxidant process [20]. Uridine was reported to reduce oxidative stress and inflammation by inhibiting the MAPK and NF- κB signaling pathways under pathological conditions [21]. Stigmasterol (**6**) is a common sterol in many plants and possesses anti-inflammatory, antioxidant, and neuroprotective activities [22]. Studies revealed that stigmasterol exhibited substantial anti-inflammatory and immune regulatory effects and has potential for the prevention and treatment of inflammation-related diseases like arthritis [23],[24]. The anti-inflammatory effect of these isolated compounds may make contributions to the activity of *n*-hexane and ethyl acetate extracts from *B. lupulina*.

4. Conclusions

Six chemical constituents including (+)-lyoniresinol (**1**), (+)-lyoniresinol 9'-*O*- β -D-glucopyranoside (**2**), 2-methyl-1,3,6-trihydroxy-9,10-anthraquinone 3-*O*- β -D-glucopyranoside (**3**), 3-*O*- β -D-glucopyranosyl (3*S*)-1-octen-3-ol (**4**), uracil (**5**), and stigmasterol (**6**) were identified from *n*-hexane and ethyl acetate

extracts of *B. lupulina* aerial parts collected in Ho Chi Minh city. To the best of our knowledge, compounds **1**, **3**, **4**, and **5** were identified for the first time from the *Barleria* genus.

References

1. Gangaram S., Naidoo Y., Dewir Y. H., El-Hendawy S. (2021), Phytochemicals and biological activities of *Barleria* (Acanthaceae), *Plants*, 11(1), 82.
2. Sawarkar H. A., Pandey A. K., Biyani K. R. (2023), Phytochemicals and therapeutic potentials of *Barleria lupulina*, *Bangladesh Journal of Pharmacology*, 18(4), 119-129.
3. Do T. L. (2004), *Vietnamese medicinal plants and materials*, Medical Publishing House, Hanoi.
4. Keomanykham O., Anh H. T., Duyen N. T., Giang T. N., Kim N. Q., Hoa T. T., Hoa L. T. P., Quang D. N. (2025), Iridoid glucosides from *Barleria lupulina* LINDL, *HNUE Journal of Science Natural Science*, 70(1), 53-60.
5. Lee S. R., Clardy J., Senger D. R., Cao S., Kim K. H. (2016), Iridoid and phenylethanoid glycosides from the aerial part of *Barleria lupulina*, *Revista Brasileira de Farmacognosia*, 26(3), 281-284.
6. Kim K. H., Park Y. J., Chung K. H., Yip M. R., Clardy J., Senger D., Cao S. (2015), Iridoid glycosides from *Barleria lupulina*, *Journal of Natural Products*, 78(2), 320-324.
7. Suba V., Murugesan T., Arunachalam G., Mandal S. C., Saha B. P. (2004), Anti-diabetic potential of *Barleria lupulina* extract in rats, *Phytomedicine*, 11(2-3), 202-205.
8. Ilori N. T. O., Liew C. X. Q., Fang C. M. (2020), The anti-inflammatory properties of *Acanthus ebracteatus*, *Barleria lupulina* and *Clinacanthus nutans*: A systematic review, *Molecular Biology Reports*, 47(12), 9883-9894.
9. Kumari R., Kumar S., Kumar A., Goel K. K., Dubey R. C. (2017), Antibacterial, antioxidant and immuno-modulatory properties in extracts of *Barleria lupulina* Lindl, *BMC Complementary and Alternative Medicine*, 17(1), 484.
10. Nguyen N. T., Dang P. H., Nguyen H. X., Le T. H., Do T. N. V., Nguyen M. T. T. (2022), A new lignan from the stems of *Buchanania lucida* Blume (Anacardiaceae), *Natural Product Research*, 36(14), 3737-3740.
11. Pham H. Y., Dan T. T. H., Nguyen H. H., Ngo A. B., Duong T. H. Y., Nguyen A. H., Nguyen V. T., Bui H. T., Nguyen X. N., Phan V. K. (2022), Lignans from *Gymnema sylvestre*, *VNU Journal of Science: Natural Sciences and Technology*, 38(4), 18-24.
12. Yuasa K., Ide T., Otsuka H., Ogimi C., Hirata E., Takushi A., Takeda Y. (1997), Lignan and neolignan glycosides from stems of *Alangium premnifolium*, *Phytochemistry*, 45(3), 611-615.
13. Itokawa H., Qiao Y., Takeya K. (1989), Anthraquinones and naphthohydroquinones from *Rubia cordifolia*, *Phytochemistry*, 28(12), 3465-3468.
14. Yamamura S., Ozawa K., Ohtani K., Kasai R., Yamasaki K. (1998), Antihistaminic flavones and aliphatic glycosides from *Mentha spicata*, *Phytochemistry*, 48(1), 131-136.
15. Ha N. T. T., Tra N. T., Ninh B. H., Tiep N. K. (2022), Chemical constituents from the stem barks of *Polyscias serrata* Balf, *Vietnam Journal of Science and Technology*, 60(6), 966-973.
16. Khoi N. M., Luc T. M. (2016), Isolation of stigmaterol from methanolic extract of stem and foliage of *Abrus precatorius* Linn, *Journal of Science and Technology, the University of Danang*, 12(109), 39-42.
17. Guo X. H., Pang L., Gao C. Y., Meng F. L., Jin W. (2023), Lyoniresinol attenuates cerebral ischemic stroke injury in MCAO rat based on oxidative stress suppression via regulation of Akt/GSK-3 β /Nrf2 signaling, *Biomedicine & Pharmacotherapy*, 167, 115543.
18. Lu C. L., Zhu W., Wang D. M., Chen W. L., Hu M. M., Wang M., Lu C. J. (2015), Inhibitory effects of chemical compounds isolated from the rhizome of *Smilax glabra* on nitric oxide and tumor necrosis factor- α production in lipopolysaccharide-induced RAW264.7 cell, *Evidence-Based Complementary and Alternative Medicine*, 2015(1), 602425.
19. Lee D. G., Jung H. J., Woo E. R. (2025), Antimicrobial property of (+)-lyoniresinol-3 α -O- β -D-glucopyranoside isolated from the root bark of *Lycium chinense* Miller against human pathogenic microorganisms, *Archives of Pharmacal Research*, 28, 1031-1036.
20. Yang Y., Ye Y., Deng Y., Gao L. (2024), Uridine and its role in metabolic diseases, tumors, and neurodegenerative diseases, *Frontiers in Physiology*, 15, 1360891.
21. Luo Y., Chen H., Huang R., Wu Q., Li Y., He Y. (2021), Guanosine and uridine alleviate airway inflammation via inhibition of the MAPK and NF- κ B signals in OVA-induced asthmatic mice, *Pulmonary Pharmacology & Therapeutics*, 69, 102049.
22. Li J., Zheng X., Qi J. (2025), Research progress on the therapeutic mechanisms of stigmaterol for multiple diseases, *Molecules*, 30(9), 1874.
23. Xie W., Yang H., Guo C., Xie R., Yu G., Li Y. (2023), Integrated network pharmacology and experimental validation approach to investigate the mechanisms of stigmaterol in the treatment of rheumatoid arthritis, *Drug Design, Development and Therapy*, 17, 691-706.
24. Ahmad-Khan M., Sarwar A., Rahat R., Ahmed R. S., Umar S. (2020), Stigmaterol protects rats from collagen induced arthritis by inhibiting proinflammatory cytokines, *International Immunopharmacology*, 85, 106642.